

# An Investigation on the Mathematical Models for the Growth of Some Tumor Cells and Their Control

Manish Sharma<sup>1\*</sup>, Sanjay Kumar Pandey<sup>2</sup>, Shailendra Kumar<sup>3</sup>

<sup>1</sup>Assistant Professor, Department of Mathematics, Shri L. B. S. Degree College, Gonda (India)
<sup>2</sup>Professor, Department of Mathematics, Shri L. B. S. Degree College, Gonda (India)
<sup>3</sup>Assistant Professor, Department of Mathematics, Brahmanand College, Kanpur, U.P.
\*Email: manishtsharma12@gmail.com

## Abstract:

Theoretical, experimental, and clinical methods for comprehending the dynamics of cancer cells and their interactions with the immune system have undergone major advancements over the past few decades. They have sparked the creation of crucial cancer treatment techniques as biotherapy, immunotherapy, chemotherapy, targeted medication therapy, and several others. Additionally, there have been notable advancements in computational and analytic models to aid in shedding light on clinical data. Our findings seem to indicate that the model used is a solid choice for researching tumour cell dynamics and that it aids in presenting the active interactions amid tumour cells, the immune system, and drug response systems.

Keywords: Mathematical, Tumor, Cancer

# 1. Introduction

Mathematical modelling allows for the simulation of the dynamics of complicated systems, which can then be used to test hypotheses and conduct experiments. In addition to shedding light on the molecular foundations of dynamical systems, scientific models may be used to simulate these systems in a short length of time, saving researchers the costly costs of lab trials and the associated lifestyle adjustments. Such replicas dismiss be standardized using new or clinical statistics, and before clinical intervention, conflicting hypotheses of tumour progression can be thoroughly examined. This is especially true for oncology. There are several quantitative modeling methods, and more and more theoretical methods are being effectively applied to malignant growth science. Individual cell models and differential condition models prepared for quantitative malignant growth science more than twenty years ago. Here, we will provide an overview of how these models are created and how to use them to mimic tumour development and therapeutic response. Afterwards, we'll talk about a variety of models and their accuracy in predicting and confirming cancer biology. Bodnar et. al. (2016) developed a set of models for angiogenesis that generalise those of Hahnfeldt et al. Two differential equations with random time delays are used to represent the considered model family. It is shown that the answers exist and are unique on a global scale. 2021, Gabriella Bretti et al. The recruitment of immune cells to a tumour is an important predictor of prognosis and therapeutic response in cancer, and the complex interplay between cells, extracellular molecules, and secreted chemotactic factors is

crucial in regulating the migration of both activating and suppressive immune cell types to the site of disease.

#### 2. ODE models of tumor growth

Because time is constantly changing, it is challenging to estimation how countless cancer cells are present in a tumour. Tumor cells can grow, remain dormant, or pass away. This makes recitation the amount of tumour cells as a meaning of period incredibly difficult. Yet, formalizing the projected changes in cell number as time passes is simple. Individual when cells divide or decease do the number of breathing cells modification: The pace at which cells divide and multiply depends on the time interval being considered, or dt. Let's don that a random cancer cell has a 24-hour cell cycle. The likelihood that the cell will divide over the course of one day is then very likely to be 100%. Without knowing where a cell is in the cell cycle right now, we can infer that the likelihood that it will divide within an hour is 1/24. The aforementioned illustration can be unswervingly applied to the inhabitant's neck and neck despite short of eloquent the precise integer of cells in a tumour inhabitants. Similar to this, just a portion of the inhabitants cells (about 1/24) are anticipated to division if dt=1 hour. Similar explanations exist for prison cell death. We so requirement presents the period transformation as healthy as binary limits addicted to the upstairs equation:

 $\frac{\text{difference in quantity of cells}}{dt} = +\alpha \text{ (quantity of cells)} - \beta \text{ (quantity of cells)},$ 

Where  $\alpha$  and  $\beta$  signify the apiece capita development and death charges of the entire cell inhabitants and are, individually, unspoken as the segment of separating and failing cells both dt. It goes without saying that the number of cells must rise following a proliferative happening and fall following a cell death episode. Familiarize the c variable, which stands for cells. The equation above can be expressed as follows using the modification in cell integer as the variable dc:

$$\frac{dc}{dt} = \alpha c - \beta c \tag{1}$$

Such equivalence is baptized an ODE. Occupancy us undertake that at period *t*=0, the initial opinion of a trial, we must single million prison cell, i.e.  $c=10^6$ . Populace progress subtleties can shadow unique of 3 fortunes: (i) if  $\alpha = \beta$ , before dc/dt = 0. In this circumstance the integer of compartments in the inhabitants fixes not variation plus the inhabitants exhibitions a government of growth latency. It stays of letter that whichever  $\alpha = \beta = 0$ , that is entirely cubicles in everybody are in common of cell inexpression, or  $\alpha = \beta > 0$  in which event cell explosion is composed by booth decease, If (ii)  $\alpha > \beta$ , and the cell populace resolve incessantly breed thru bigger  $\alpha$ - $\beta$  rates springy quicker progress. Scheduled the supplementary pointer, the inhabitants resolve monotonically decline if (iii)  $\alpha < \beta$  and hence dc/dt < 0 (fig. 1).



Fig. 1: Development subtleties of cell inhabitants *c* over period *t* on behalf of diverse comparative rates of cell propagation  $\alpha$  and booth passing  $\beta$ ;  $c = 10^6$  cells at period *t*=0.

Eqn. (1) canister be dense to a one-boundary problematic. The footings  $\alpha c \cdot \beta c$  dismiss be shared into the solitary term  $(\alpha \cdot \beta)c$ , and we present the solitary stricture,  $\lambda$ ,  $\lambda = \alpha \cdot \beta$ , which is baptized the net populace growing rate. The PDE recitation cell inhabitant's revolution over period is then

$$\frac{dC}{dt} = \lambda c \tag{2}$$

For example earlier, if  $\lambda < 0$ ,  $\lambda = 0$ , or  $\lambda > 0$  the inhabitants declines, leftovers at a endless size, or surges correspondingly. New statistics after *in vitro* or *vivo* inhabitant's educations dismiss then is castoff to parameterize such prototypical (fig. 2).



Fig. 2. A) Mock inhabitants' growing analogous to *in vitro* new statistics with 5% average error saloons and designed leaning line. B) Precise prototypical consequences of inhabitant's progress for diverse strictures of  $\lambda$ .

## 3. Dynamic tumor development rates

In order to simulate tumour growth, a continuous apiece capita evolution rate is defined, which results in exponentially expanding cell populations while ignoring carrying capacity restrictions. Solid tumours develop quickly at first but slow down as they get bigger. As a result, rather than being constant, the apiece capita rate of tumour progress should be dependent on the tumour size.

$$\frac{dC}{dt} = f(c)c$$

A typical instance of a apiece capita proportion of lump development reliant on tumor magnitude comparative to the host loud size *K* is agreed by logistic perfect.

$$\frac{dc}{dt} = \lambda c \left(\frac{K-c}{K}\right) = \lambda c \left(1 - \frac{c}{K}\right) \tag{3}$$

Now  $f(c) = \left(1 - \frac{c}{\kappa}\right)$  and consequently the apiece capita advancement rate cuts as c increases. Assuming that cancer size c << K,  $before\left(1 - \frac{c}{\kappa}\right) \rightarrow 0$  and inhabitants development is hindered. Fig. 3 expressions the development of people magnitude as fine as the developing rate subject to occupants size outright to K for unlike constraints  $\lambda$ .



Fig. 3: Inhabitants growing imperfect by resonant volume K. A) Inhabitants size as afunction of

period with eqn. (2) B) Conforminsg active each capita development rates.

There have been many discussions about dynamic growing rate purposes that are relevant to tumour development. It has remained demonstrated that the so-called Gompertz curve can mimic biological growing that slows with inhabitant's size, making it applicable to the observed decrease in tumour development with tumour size. The growth rate is calculated as the carrying capacity divided by the negative logarithm of the recent inhabitants' size:

$$\frac{dC}{dt} = -\lambda c \, \log(1 - c/K) \, (4)$$

Notwithstanding an additional rapid development degree fall-off, inhabitants development impersonators that experiential with eqn. (3) and expected in fig. 3 (fig. 4).



**Fig. 4. Gompertz inhabitants development restricted by resonant size** *K***. A)** Populace size as a meaning of period by eqn. (4). B) Conforming dynamic apiece capita development charges.

The resounding capacity may not be continual in a diseased environment. Depending on the size of the tumour, an upsurge in cell physique and physical gravity can result in the distortion and development of the cellar membrane. Yet, tumour growth can also slow in the absence of vital nutrients and growth hormones supplied by the swarm vasculature. The transport of oxygen besides nutrients obsessed by the centre of the tumour is constrained during vascular dormancy, resultant in poise between cell circulation and cell passing. The largest tumour extent that dismiss be maintained by circulatory stock is then determined by carrying capacity. The tumour inhabitants cooperates by the swarm vasculature by releasing pro- and anti-antigenic factors, which results in a resonant bulk K that varies over period. Finished diffusion feeding influences Hahnfeldt resulting that vasculature hang-up is comparative to the (cancer volume)<sup>2/3</sup>. In its place of being continuous, vicissitudes in *K* are spoken by

$$\frac{dK}{dt} = \emptyset c - \emptyset K c^{2/3}$$
(5)

wherever  $\phi$  besides  $\varphi$  stand endless optimistic charges of angiogenesis motivation and shyness, correspondingly. Primarily lump evolution is quicker payable to plentiful stimulus nevertheless as the cancer cultivates the inhibitory properties motivation compensate the stimulator elastic a mesa vascular hoard and growth dimension (fig. 5).



Fig. 5. Tumor growth by a go-ahead resonant capacity A) Fruition of tumor extent *c* and booming size *K* thru  $\lambda$ =0.182,  $\phi$ =5.75, and  $\varphi$ =0.00773, restrictions formfitting to Lewis lung carcinoma vertical in C56BL/6 mice.B) Conforming dynamic apiece capita development rates.

#### 4. Modeling cancer treatment

There are two methods for treating tumours. By causing cancer cells to die while they are still dividing or by reducing the carrying capacity, one can significantly reduce the size of the tumour. Differential equation models may easily include the impacts of both cancer therapy options (eqns. (4) & (5)). The most basic form of anti-tumor therapy results in a constant tumour cell kill by an asset of  $0 \le \xi \le 1$ , comparable to chemotherap. Fig. 6 depicts the cancer's reaction to such cubicle killing at various intensities afterward the tumour has full-grown aimed at 39 days (c.f. Fig. 5). The inhibition of angiogenesis caused by anti-antigenic medicines administered was hypothesized by Hahnfeldt and colleagues to be comparative to the medication consideration g(t), which contains not completely empty medication focuses after prior directions.

$$g(t) = \int_0^t a(t') \exp\left(-clr(t-t')\right) dt',$$

everywhere a(t') besides clr(t-t') stand the charges at which the inhibitor thought is coordinated and its endorsement rate, individually.



Fig. 6. Anti-tumor action for dissimilar asset cell kills  $\xi$ .

#### 5. Partial differential equation (PDE) representations of tumor growth

The most obvious drawback of this technique is the absence of spatial thought, despite the fact that ODE models obligate shown to be a valuable utensil for simulating the growth of the overall tumour cell number concluded period. The initial tumours locally infiltrate the nerve and meal to reserved regions of the bulk to produce subordinate tumours, not because there are overall more cancer cells in the patient's body. The primary factor in cancer patients' deaths is these metastatic tumours. PDE models can be used to mimic the spatially based processes of cancer invasion and metastatic dissemination. In these models, the number of residents n at any given longitudinal point (x), (x, y), or (x, y, z) in one, two, or three aspects, separately, is many times expressed as a mass, or a negligible part of the greatest open volume at this spot, and is scaled somewhere in the range of 0% and 100 percent, or 0 and 1. The variable n is now dependant on vicissitudes in the considered spatial dimensions as well as changes in time (t). Hence, the partial derivatives of its derivatives are required in the equation for n. overlooking the measured 3-D sphere, the PDE of *n* w.r.t *t* is inscribed as  $\frac{\partial n}{\partial t}$ .

The key to the development and spread of cancer is tissue invasion, which is essential for effective metastasis. Three main elements make up the invasion process: The extracellular matrix (ECM) or surrounding tissue is destroyed by the numerous matrix degrading enzymes (MDEs) discharged by the malignant growth cells. Additionally, the danger cells forcefully attack the nearby tissue through immigration and spread. Our model took into account how too many H+ ions could cause local tissue to deteriorate, creating a place where cancer cells could spread and grow. The extracellular atmosphere, H+ ions, and the set of PDE secondhand to reenact the spatiotemporal fulfillment of growth cell c, are as per the following:

$$\frac{\partial c}{\partial t} = \nabla . \left( D_c (1 - v) \nabla_c \right) + \rho c (1 - c)$$
$$\frac{\partial m}{\partial t} = \nabla^2 m + \delta (c - m)$$
$$\frac{\partial v}{\partial t} = \nabla . \left( (1 - v) - \eta m v \right)$$

where Dc is a constant value, is a constant value for the rate at which cancer cells divide, is a constant value for the rate at which H+ ions are created (which is believed to be equivalent to the rot rate), and is a consistent incentive for the rate at which the extracellular grid debases. The previously mentioned conditions outline how disease cells multiply, go through nonlinear dispersion (which shifts relying upon the thickness of typical tissue; high thickness of ordinary tissue brings about lower dissemination, while low thickness of ordinary tissue brings about lower dissemination, while low thickness of ordinary tissue brings about lower dissemination, while low thickness of ordinary tissue brings about lower dissemination, while low thickness of ordinary tissue. In the Gatenby and Gawlinski model, the development and rot of H+ particles are accepted to continue at a similar rate for mathematical straightforwardness, permitting the ordinary tissue to accomplish its sound state without even a trace of disease cells by linear decay. The mathematical formulation of diffusion, the population-level description of random cell movement, is outside the extent of this article, but the intrigued peruser is alluded to course books on halfway differential conditions. Utilizing a combination of virtual experience and numerical examination (voyaging wave hypothesis) of the previously mentioned arrangement of fractional differential conditions, this was thusly identified in hematoxylin-eosin (H&E) stained micrographs of invasive tumours (Fig. 7).



Figure 7: Experimental validation of the cancer invasion and simulation findings.

Later PDE models expanded on this concept and used reaction-diffusion-taxis models to study the interactions amid cancer cells (c), tissue (v), and debasing enzymes (m), where the development of haptotaxis frolicked a crucial part in the immigration of the malignancy cells. Particular attention was paid to the function of haptotaxis in Anderson's work, which made use of a two-dimensional context for the first time and clearly recognised the effect of the tissue.

$$\frac{\partial c}{\partial t} = D_c \nabla^2 c - \eta \nabla (\mathbf{c} \nabla \mathbf{v})$$
$$\frac{\partial m}{\partial t} = \nabla^2 m + \delta(c - m)$$
$$\frac{\partial v}{\partial t} = -\eta m v$$

The band model did not place all the weight on the main character of cancer cell migration in its incursion, but rather on cell production as well.

## 6. Discrete models of tumor growth

Anderson paper stayed the first to yield into account a distinct model of malignancy cell invasion, consequential since the continuum PDE model, in accumulation to presence the initial 2-layered range exemplary of disease intrusion fixing with respect to haptotaxis. The computational recreation discoveries of this model examined the finding that a surgeon-detectable "visible margin" of malignant tissue can be crossed by a single cancer cell. This study may have been the first to examine the relationship between probability and stochastic events in invasion models. A taster outcome of a calculation multiplication of the different model in a 2-layered area is shown in Figure 8.





The ability to consider actions at the close of individual cells is one benefit discrete replicas have finished continuum models. Important occurrences like mutations as well as various phenotypic traits can be considered using discrete models. The creation of "multistate models," everyplace intracellular processes dismiss be described via organisms of ODE and then allied to cellular level characteristics, has also been facilitated by the introduction of discrete models. For the past 15 years or so, cross continuum-discrete mockups need too been created to mimic (tumor-induced) angiogenesis, another crucial stage in the growth

and evolution of solid tumours. In their seminal study, Anderson and Chaplain first proposed the following organization of PDEs to model the spatiotemporal connections among (tip) n, extracellular atmosphere, v, and tumour antigenic factor (TAF). In addition to directed passage via chemo taxis in reaction to TAF slopes (such as VEGF), the endothelial cells also engage in random walking plus haptotaxis, which is consumed relocation in answer to fibronectin inclines in the extracellular atmosphere. As shown in Figure 9, the computational reproduction outcomes since the disconnected kind of the typical that included cellular androids like activities, such as vessel diverging and anastomosis, stayed clever to replicate pronged, connected vessel grids as conveyed in trial reports. Today, such discrete models have been created that take into account the impact of blood run in the interior vessels. There are already a number of complimentary models that focus on distinct facets related to tumour angiogenesis and the development of vascular tumours.



Figure 9: The simulated results of a unique angiogenesis model.

Gradients of TAF and fibronectin cause discrete tubes to react chemo tactically besides haptotactically, forming a split, linked capillary system that container interact by the solid tumour. Moreover, these separate angiogenesis mockups have stood extended and established to mimic added classifications where the evolution of family vessels dramas a crucial heroine, such as wound medicinal and retinal change.

### 7. Discussion

In the last couple of decades, cancer research has benefited from an avalanche of mathematical models. Within this text, we must provide an example of how complex measureable replicas remain industrialized and likened by new statistics, and presented in what way they container be rummage-sale to fake multifaceted organic developments and connections. We must selected influential IDs as instances, and aimed at ease must had to permission out a big form of brilliant precise showing literature. The reader who is really interested in the current status of cancer research will be referred to recent evaluations of books and articles that provide a fuller picture of the field.

#### **References:**

- 1. Orme, M. E., & Chaplain, M. A. J. (1997). Two-dimensional models of tumour angiogenesis and antiangiogenesis strategies. *Mathematical Medicine and Biology: A Journal of the IMA*, *14*(3), 189-205.
- Ucar, E., Özdemir, N., & Altun, E. (2019). Fractional order model of immune cells influenced by cancer cells. *Mathematical Modelling of Natural Phenomena*, 14(3), 308.
- Vaghi, C., Rodallec, A., Fanciullino, R., Ciccolini, J., Mochel, J. P., Mastri, M., ... & Benzekry, S. (2020). Population modeling of tumor growth curves and the reduced Gompertz model improve prediction of the age of experimental tumors. *PLoS computational biology*, *16*(2), e1007178.
- Ruan, S. and Wei, J., 2003. On the zeros of transcendental functions with applications to stability of delay differential equations with two delays. Dynamics of Continuous Discrete and Impulsive Systems Series A, 10, pp.863-874.
- 2. Xu, J., Vilanova, G. and Gomez, H., 2016. A mathematical model coupling tumor growth and angiogenesis. PloS one, 11(2), p.e0149422.s
- Ruan, S, Wang, W: Dynamical behavior of an epidemic model with a nonlinear incidence rate. J. Differ. Equ. 188, 135-163 (2003)
- Xiao, D, Ruan, S: Global analysis of an epidemic model with nonmonotone incidence rate. Math. Biosci. 208, 419-429 (2007)
- 5. Yang, Y, Xiao, D: Influence of latent period and nonlinear incidence rate on the dynamics of SIRS epidemiological models. Discrete Contin. Dyn. Syst., Ser. B 13, 195-211 (2010)s
- 6. De Pillis, LG, Radunskaya AE and Wiseman CL,. "A validated mathematical model of cell-mediated immune response to tumor growth". Cancer research, 65(17), (2005), pp.7950-7958.
- 7. Fry, B., 2009. A mathematical model of diffusion-driven tumor growth with viral therapy.
- Ghosh, D., Khajanchi, S., Mangiarotti, S., Denis, F., Dana, S.K. and Letellier, C., 2017. How tumor growth can be influenced by delayed interactions between cancer cells and the microenvironment?. *BioSystems*, 158, pp.17-30.
- Bodnar M and Piotrowska M J. "Stability analysis of the family of tumour angiogenesis models with distributed time delays". Commun Nonlinear Sci Numer Simulat. (2016), pp. 124–142.
- Mahardika, R., & Sumanto, Y. D. (2019, May). Routh-hurwitz criterion and bifurcation method for stability analysis of tuberculosis transmission model. In *Journal of Physics: Conference Series* (Vol. 1217, No. 1, p. 012056). IOP Publishing.
- Peirce S M. "Computational and Mathematical Modeling of Angiogenesis". Microcirculation (2008). 15(8). pp.739–751.
- 12. Kuang, Y: Delay Differential Equations with Applications in Population Dynamics. Academic Press, San Diego (1993)

- 13. van der Driessche, P, Watmough, J: Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. Math. Biosci. 180, 29-48 (2002)
- 14. Brauer, F, van der Driessche, P, Wu, J: Mathematical Epidemiology. Lecture Notes in Mathematics, vol. 1945. Springer, New York (2008)
- 15. Zhang, ZF, Ding, TR, Huang, WZ, Dong, ZX: Qualitative Theory of Differential Equations. Translations of Mathematical Monographs, vol. 101. Am. Math. Soc., Providence (1992)
- 16. Lasalle, JP: An invariance principle in the theory of stability. In: International Symposium on Differential Equations and Dynamical Systems, pp. 277-286. Academic Press, New York (1967)

# Cite this Article:

Manish Sharma, Sanjay Kumar Pandey, Shailendra Kumar, "An investigation on the mathematical models for the growth of some tumor cells and their control ", International Journal of Scientific Research in Modern Science and Technology (IJSRMST), ISSN: 2583-7605 (Online), Volume 2, Issue 5, pp. 22-32, May 2023. Journal URL: <u>https://ijsrmst.com/</u>